

Supporting Information

Porter et al. 10.1073/pnas.0808010105

SI Text

Effect of Variation of [CheA₃] on the Dephosphorylation Rate of CheY₆-P. The dephosphorylation rate of CheY₆-P was measured in the presence of varying amounts of CheA₃ (Fig. S1). As the concentration of CheA₃ was increased, the rate of CheY₆-P dephosphorylation increased, with no evidence of saturation over the concentration range tested. The limited solubility of CheA₃ coupled with the high concentrations of CheY and phosphodonors needed for this assay, imposed the upper limit of the CheA₃ concentration range tested in these assays of 2.5 μM. The shape of the curve is suggestive of sigmoidal kinetics with respect to CheA₃, possibly indicating either positive cooperativity or some need for CheA₃ to oligomerize to function as a phosphatase. Either way, these data suggest that higher concentrations of CheA₃ might be able to stimulate CheY₆-P dephosphorylation by more than the factor of 3 seen for 2.5 μM CheA₃.

CheA₃ Is Not a Homologue of the CheC Family of Phosphatases. Sequence analysis of the 794-amino acid region between the P1 and P5 domain of CheA₃ revealed a partial match (D-X₄-E-X₂-

N-X₂₀-P) to the consensus sequence of the CheC family of CheY-P phosphatases (D/S-X₃-E-X₂-N-X_{21/22}-P) (1, 2). The E and N residues within this consensus sequence have been shown to be essential for phosphatase activity in a number of CheC family members (2, 3). To determine whether CheA₃ uses a similar phosphatase mechanism, the corresponding residues in CheA₃ (E585 and N588) were both changed to serine and the resulting protein assayed for CheY₆-P phosphatase activity. The CheA₃(E585S,N588S) protein displayed wild-type levels of CheY₆-P phosphatase activity (Fig. 3) indicating that CheA₃ is not a member of the CheC family of phosphatases.

CheA₃ Does Not Show Phosphatase Activity Toward CheA₂-P or CheA₃P1-P. Experiments where CheA₃ was incubated alone with the phosphodonors, CheA₂-P and CheA₃P1-P, in the absence of added RRs, detected no change in the concentration of phosphodonor over time (Fig. S2), showing that CheA₃ does not possess histidyl-phosphate phosphatase activity toward either of the phosphodonors used in this study.

1. Szurmant H, Bunn MW, Cannistraro VJ, Ordal GW (2003) *Bacillus subtilis* hydrolyzes CheY-P at the location of its action: The flagellar switch. *J Biol Chem* 278:48611–48616.
2. Muff TJ, Ordal GW (2007) The CheC phosphatase regulates chemotactic adaptation through CheD. *J Biol Chem* 282:34120–34128.
3. Park SY, et al. (2004) Structure and function of an unusual family of protein phosphatases: The bacterial chemotaxis proteins CheC and CheX. *Mol Cell* 16:563–574.
4. Penfold RJ, Pemberton JM (1992) An improved suicide vector for construction of chromosomal insertion mutations in bacteria. *Gene* 118:145–146.
5. Sockett RE, Foster JCA, Armitage JP (1990) Molecular biology of the *Rhodobacter sphaeroides* flagellum. *FEMS Symp* 53:473–479.
6. Porter SL, Armitage JP (2004) Chemotaxis in *Rhodobacter sphaeroides* requires an atypical histidine protein kinase. *J Biol Chem* 279:54573–54580.
7. Porter SL, et al. (2006) The CheYs of *Rhodobacter sphaeroides*. *J Biol Chem* 281:32694–32704.
8. Porter SL, Warren AV, Martin AC, Armitage JP (2002) The third chemotaxis locus of *Rhodobacter sphaeroides* is essential for chemotaxis. *Mol Microbiol* 46:1081–1094.
9. Wadhams GH, Warren AV, Martin AC, Armitage JP (2003) Targeting of two signal transduction pathways to different regions of the bacterial cell. *Mol Microbiol* 50:763–770.
10. Shah DS, et al. (2000) Identification of a fourth cheY gene in *Rhodobacter sphaeroides* and interspecies interaction within the bacterial chemotaxis signal transduction pathway. *Mol Microbiol* 35:101–112.
11. Porter SL, Armitage JP (2002) Phosphotransfer in *Rhodobacter sphaeroides* chemotaxis. *J Mol Biol* 324:35–45.
12. Martin AC, et al. (2001) CheR- and CheB-dependent chemosensory adaptation system of *Rhodobacter sphaeroides*. *J Bacteriol* 183:7135–7144.
13. Schäfer A, et al. (1994) Small mobilizable multipurpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19 - Selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145:69–73.




Fig. S1. The effect of varying the concentration of CheA₃ on the rate of CheY₆-P dephosphorylation. Reactions contained 30 μM CheA₃P1-P, 400 μM CheY₆, and the concentration of CheA₃ was varied between 0 μM and 2.5 μM. Error bars show standard error of the mean obtained from six replicates.




Fig. S2. Phosphorimages of SDS/PAGE gels showing the effect of CheA₃ on levels of CheA₂-P and CheA₃P1-P. (A) 2 μM CheA₂-P was incubated in the absence (left half of gel) and presence of 2.5 μM CheA₃ (right half of gel). (B) 2 μM CheA₃P1-P was incubated in the absence (left half of gel) and presence of 2.5 μM CheA₃ (right half of gel). Ten-μl reaction samples were taken at the time points indicated and quenched in 20 μl of 1.5× SDS/EDTA loading dye. The quenched samples were analyzed by SDS/PAGE and detected by phosphorimaging.

